arXiv:1602.02577v1 [math.GR] 8 Feb 2016

HEXATONIC SYSTEMS AND DUAL GROUPS IN MATHEMATICAL MUSIC THEORY

CAMERON BERRY AND THOMAS M. FIORE

Abstract. Motivated by the music-theoretical work of Richard Cohn and David Clampitt on late-nineteenth century harmony, we mathematically prove that the PL-group of a hexatonic cycle is dual (in the sense of Lewin) to its T/I-stabilizer. Our point of departure is Cohn's notions of maximal smoothness and hexatonic cycle, and the symmetry group of the 12-gon; we do *not* make use of the duality between the T/I-group and PLR-group. We also discuss how some ideas in the present paper could be used in the proof of T/I-PLR duality by Crans–Fiore–Satyendra in [4].

1. Introduction: Hexatonic Cycles and Associated Dual Groups

Why did late nineteenth century composers, such as Frank, Liszt, Mahler, and Wagner, continue to privilege consonant triads over other tone collections, while simultaneously moving away from the diatonic scale and classical tonality?

Richard Cohn proposes an answer in [3], independent of acoustic consonance: major and minor triads are preferred because they can form *maximally smooth cycles*. Consider for instance the following sequence of consonant triads, called a *hexatonic cycle* by Cohn.

$$E\flat, e\flat, B, b, G, g, E\flat \tag{1}$$

We have indicated major chords with capital letters and minor chords with lowercase letters. Although the motion from a major chord to its parallel minor, e.g. $E\flat$ to $e\flat$, B to b, and G to g, is distinctly non-diatonic, this sequence has cogent properties of importance to late-Romantic composers, as axiomatized in Cohn's notion of maximally smooth cycle [3, page 15].

- It is a *cycle* in the sense that the first and last chords are the same but all others are different. A *cycle* is required to contain more than three chords.
- All of the chords are in one "set class," in this case each chord is a consonant triad.
- Every transition is *maximally smooth* in the sense that two notes stay the same while the third moves by the smallest possible interval: a semitone.

Cohn considered movement along this sequence transformationally as an action by a cyclic group of order 6. Additionally, David Clampitt considered in [2] movement along this sequence via P and L, and also via certain rotations and reflections. As usual, we denote by P the "parallel" transformation that sends a major or minor chord to its parallel minor or major chord, respectively. We denote by L the "leading tone exchange" transformation, which moves the root of a major chord down a semitone and the fifth of a minor chord up a semitone, so the L sends consonant triads $e\flat$ to B,

and b to G, and g to $E\flat$. The hexatonic cycle (1) is then positioned in network (2), with alternating P and L transformations between the nodes.

(2)


Wagner's Grail motive in *Parsifal* can be interpreted in terms of the network (2), as proposed by David Clampitt in [2]. A small part of Clampitt's analysis of the first four chords is pictured in Figure 1. Clampitt includes the final $D\flat$ chord, which lies outside of the hexatonic cycle (1), into his interpretation via a conjugation-modulation applied to a certain subsystem. A third interpretation, in addition to the cyclic one of Cohn [3, Example 5] and the PL interpretation in Figure 1, was also proposed by Clampitt, this time in terms of the transpositions and inversions $\{T_0, T_4, T_8, I_1, I_5, I_9\}$. Clampitt observes that this group and the PL-group form *dual groups in the sense of Lewin* [14], via their actions on the hexatonic set of chords in (1). The perceptual basis of all three groups are explained in [2].

$E\flat$, b, G, $e\flat$, $D\flat$

(A) Chord sequence in Grail motive from Wagner, *Parsifal*, Act 3, measures 1098–1100, see Clampitt [2, Example 1].



(B) First four chords of the Grail motive in a hexatonic PL-network of Clampitt. Notice that the bottom arrow is the composite of the three top arrows, and goes in the opposite direction of the bottom arrow of diagram (2).

FIGURE 1. Wagner's Grail motive and Clampitt's interpretation of first four chords using the PL-network in diagram (2).

The contribution of the present article is to directly prove that the PL-group and the group $\{T_0, T_4, T_8, I_1, I_5, I_9\}$ in Clampitt's article are dual groups acting on (1). Our point of departure is the hexatonic cycle (1), the standard action of the dihedral group of order 24 on the 12-gon, and the Orbit-Stabilizer Theorem. We do *not* use the duality of the T/I-group and PLR-group. Some arguments in Section 3 are similar to arguments of Crans–Fiore–Satyendra [4], but there are important differences, see Remark 3.9.

Just how special are the consonant triads with regard to the maximal smoothness property? According to [3], only 6 categories of tone collections support maximally smooth cycles: singletons, consonant triads, pentatonic sets, diatonic sets, complements of consonant triads, and 11-note sets. Clearly the singletons and 11-note sets do not give musically significant cycles. The pentatonic sets and the diatonic sets each support only one long cycle, which exhausts all 12 of their respective exemplars. The consonant triads and their complements, *on the other hand,* support short cycles that do not exhaust all of their transpositions and inversions. The maximally smooth cycles

of consonant triads are enumerated as sets as follows.

$$\{E\flat, e\flat, B, b, G, g\} \tag{3}$$

$$\{E, e, C, c, A\flat, a\flat\} \tag{4}$$

$$\{F, f, C\sharp, c\sharp, A, a\} \tag{5}$$

$$\{F\sharp, f\sharp, D, d, B\flat, b\flat\} \tag{6}$$

These are the four *hexatonic cycles* of Cohn, see [3, page 17]. They (and their reverses) are the only short maximally smooth cycles that exist in the Western chromatic scale.

2. Mathematical and Musical Preliminaries: Standard Dihedral Group Action on Consonant Triads and the Orbit-Stabilizer Theorem

We quickly recall the standard preliminaries about consonant triads, transposition, inversion, P, L, and the Orbit-Stabilizer Theorem. A good introduction to this very well-known background material is [4]. Since this background has been treated in many places, we merely rapidly introduce the notation and indicate a few sources.

2.1. **Consonant Triads.** We encode pitch classes using the standard $\mathbb{Z}_{12}$ model, where $C = 0$, $C\sharp = D\flat = 1$, and so on up to $B = 11$. Via this bijection we freely refer to elements of $\mathbb{Z}_{12}$ as *pitch classes*. Major chords are indicated as ordered 3-tuples in $\mathbb{Z}_{12}$ of the form $\langle x, x+4, x+7\rangle$, where x ranges through $\mathbb{Z}_{12}$. Minor chords are indicated as 3-tuples $\langle x+7, x+3, x\rangle$ with $x \in \mathbb{Z}_{12}$. We choose these orderings to make simple formulas for P and L, this is not a restriction for applications, as the framework was extended in [7] to allow any orderings. We call the set of 24 major and minor triads *Triads*, this is the set of *consonant triads*. The letter names are indicated in Table 1.

2.2. **Transposition and Inversion, and P and L.** The twelve-tone operations *transposition* $T_n\colon \mathbb{Z}_{12} \to \mathbb{Z}_{12}$ and *inversion* $I_n\colon \mathbb{Z}_{12} \to \mathbb{Z}_{12}$ are

$$T_n(x) = x + n \qquad \text{and} \qquad I_n(x) = -x + n$$

for $n \in \mathbb{Z}_{12}$. These 24 operations are the symmetries of the 12-gon, when we consider 0 through 11 as arranged on the face of a clock. In the music-theory tradition, this group is called the T/I-group (the "/" does *not* indicate any kind of quotient). The unique reflection of the 12-gon which interchanges m and n is I_{m+n}, as can be verified by direct computation.

Many composers, for instance Schoenberg, Berg, and Webern, utilized these *mod* 12 transpositions and inversions. These functions and their compositional uses have been thoroughly explored by composers, music theorists, and mathematicians, see for example Babbitt [1], Forte [10], Fripertinger–Lackner [11], Hook [12], Hook–Peck [13],

Major Triads	Minor Triads
$C = \langle 0, 4, 7\rangle$	$\langle 0, 8, 5\rangle = f$
$C\sharp = D\flat = \langle 1, 5, 8\rangle$	$\langle 1, 9, 6\rangle = f\sharp = g\flat$
$D = \langle 2, 6, 9\rangle$	$\langle 2, 10, 7\rangle = g$
$D\sharp = E\flat = \langle 3, 7, 10\rangle$	$\langle 3, 11, 8\rangle = g\sharp = a\flat$
$E = \langle 4, 8, 11\rangle$	$\langle 4, 0, 9\rangle = a$
$F = \langle 5, 9, 0\rangle$	$\langle 5, 1, 10\rangle = a\sharp = b\flat$
$F\sharp = G\flat = \langle 6, 10, 1\rangle$	$\langle 6, 2, 11\rangle = b$
$G = \langle 7, 11, 2\rangle$	$\langle 7, 3, 0\rangle = c$
$G\sharp = A\flat = \langle 8, 0, 3\rangle$	$\langle 8, 4, 1\rangle = c\sharp = d\flat$
$A = \langle 9, 1, 4\rangle$	$\langle 9, 5, 2\rangle = d$
$A\sharp = B\flat = \langle 10, 2, 5\rangle$	$\langle 10, 6, 3\rangle = d\sharp = e\flat$
$B = \langle 11, 3, 6\rangle$	$\langle 11, 7, 4\rangle = e$

TABLE 1. The set of consonant triads, denoted *Triads*, as displayed on page 483 of [4].

McCartin [18], Mead [19], Morris [20, 21, 22, 23], and Rahn [25]. Indeed, the three recent papers [11, 19, 23] together contain over 100 references.

We consider these bijective functions on $\mathbb{Z}_{12}$ also as bijective functions *Triads* $\to$ *Triads* via their componentwise evaluation on consonant triads:

$$T_n\langle x_1, x_2, x_3\rangle = \langle T_n x_1, T_n x_2, T_n x_3\rangle \qquad \text{and} \qquad I_n\langle x_1, x_2, x_3\rangle = \langle I_n x_1, I_n x_2, I_n x_3\rangle. \tag{7}$$

Also on the set *Triads* of consonant triads (with the indicated ordering), but not on the level of individual pitch classes, we have the bijective functions $P, L\colon$ *Triads* $\to$ *Triads* defined by

$$P\langle x_1, x_2, x_3\rangle = I_{x_1+x_3}\langle x_1, x_2, x_3\rangle \qquad \text{and} \qquad L\langle x_1, x_2, x_3\rangle = I_{x_2+x_3}\langle x_1, x_2, x_3\rangle. \tag{8}$$

As remarked above, P stands for "parallel" and L stands for "leading tone exchange."

We consider T_n, I_n, P, and L as elements of the symmetric group Sym(*Triads*).

Proposition 2.1. *The bijections P and L commute with T_n and I_n as elements of the symmetric group* Sym(*Triads*).

Proof. This is a straightforward computation using equations (7) and (8). This computation has been discussed in broader contexts in [8] and [9]. □

2.3. **Orbit-Stabilizer Theorem.** Suppose S is a set with a left group action by a group G. Recall that the *orbit of an element $Y \in S$* is

$$\text{orbit of } Y := \{gY \mid g \in G\}.$$

The *stabilizer group of an element $Y \in S$* is

$$G_Y := \{g \in G \mid gY = Y\}.$$

Theorem 2.2 (Orbit-Stabilizer Theorem)**.** *If a finite group G acts on a finite set S and $Y \in S$, then $|G| \,/\, |G_Y| = |\text{orbit of } Y|$.*

2.4. **Simple Transitivity.** A group action of a group G on a set S is said to be *simply transitive* if for any $Y, Z \in S$ there is a unique $g \in G$ such that $gY = Z$.

Proposition 2.3.

(1) *An action of a group G on a set S is simply transitive if and only if it is transitive and every stabilizer G_Y is trivial.*
(2) *In particular, if a finite group G acts on a set S of the same cardinality, then G acts simply transitively if and only if acts transitively, which is the case if and only if every stabilizer is trivial.*
(3) *Similarly, if a finite group G acts on a set S simply transitively, then G and S have the same cardinality.*

Proof. (1) If the action is simply transitive, then it acts transitively and for each $Y \in S$, there is only one $g \in G$ with $gY = Y$, and hence each G_Y is trivial.

Suppose G acts transitively and for every $Y \in S$, the group G_Y is trivial. Suppose $Y, Z \in S$ and $g_1, g_2 \in G$ satisfy $g_1Y = Z$ and $g_2Y = Z$. Then $Y = g_2^{-1}Z$ and $g_2^{-1}g_1Y = Y$, so $g_2^{-1}g_1 \in G_Y = \{e\}$, and finally $g_1 = g_2$.

(2) From 1 and Orbit-Stabilizer Theorem 2.2.
(3) From 1 and Orbit-Stabilizer Theorem 2.2.

□

3. Main Theorem: Hexatonic Duality

We next turn to the main result, Theorem 3.7 on Hexatonic Duality. Recall that subgroups G and H of $\mathrm{Sym}(S)$ are *dual in the sense of Lewin* [14, page 253] if each acts simply transitively on S and each is the centralizer of the other. Recall the *centralizer of G in* $\mathrm{Sym}(S)$ is

$$C(G) = \{\sigma \in \mathrm{Sym}(S) \mid \sigma g = g\sigma \text{ for all } g \in G\}.$$

Let *Hex* be the set of chords in the hexatonic cycle (1) and $\underline{Hex}$ the set of underlying pitch classes of its chords, that is,

$$Hex := \{E\flat, e\flat, B, b, G, g\},$$
$$\underline{Hex} := \{2, 3, 6, 7, 10, 11\}.$$

Our goal is to prove that the restriction of the PL-group to *Hex* and the restriction of $\{T_0, T_4, T_8, I_1, I_5, I_9\}$ to *Hex* are dual groups, and that each is dihedral of order 6. The strategy is to separately prove the unrestricted groups act simply transitively and are dihedral, and then finally to show that the restricted groups centralize each other. We begin with a characterization of the consonant triads contained in $\underline{Hex}$.

Lemma 3.1. *The only consonant triads of Table 1 contained in $\underline{Hex}$ as subsets are the elements of Hex.*

Proof. We first identify the available perfect fifths in $\underline{Hex}$ (pairs with difference 7), and then check if the corresponding major/minor thirds are in $\underline{Hex}$.

The only pairs of the form $\langle x, x+7\rangle$ are $\langle 3, 10\rangle$, $\langle 7, 2\rangle$, and $\langle 11, 6\rangle$, and we see that $x + 4$ is contained in $\underline{Hex}$ in each case, that is, 7, 11, and 3 are in $\underline{Hex}$. Thus we have the three major chords $E\flat$, G, and B, and no others.

The only pairs of the form $\langle x+7, x\rangle = \langle y, y+5\rangle$ are $\langle 2, 7\rangle$, $\langle 6, 11\rangle$, and $\langle 10, 3\rangle$, and we see that $y+8$ is contained in $\underline{Hex}$ in each case, that is, 10, 2, and 6 are in $\underline{Hex}$. Thus we have the three minor chords g, b, and $e\flat$, and no others. □

Proposition 3.2.

(1) *The only elements of the T/I-group that preserve Hex as a set are $\{T_0, T_4, T_8, I_1, I_5, I_9\}$, so they form a group denoted H.*
(2) *$H := \{T_0, T_4, T_8, I_1, I_5, I_9\}$ is dihedral of order 6.*

Proof. (1) If an element of the T/I-group preserves *Hex* as a set, than it must also preserve the collection $\underline{Hex}$ of underlying pitch classes as a set. Geometric inspection of the plot of $\underline{Hex}$ in Figure 2 reveals that the only rotations that preserve $\underline{Hex}$ are T_0, T_4, and T_8.



FIGURE 2. The solid circles represent the subset $\underline{Hex}$ of $\mathbb{Z}_{12}$. The symmetry of the subset makes apparent that the only rotations which preserve $\underline{Hex}$ are T_0, T_4, and T_8. The geometric locations of the solid circles also imply that the reflections across the dashed lines are the only reflections which preserve $\underline{Hex}$.

Again looking at Figure 2, we see that the three reflections which interchange $2 \leftrightarrow 3$, or $6 \leftrightarrow 7$, or $10 \leftrightarrow 11$ preserve $\underline{Hex}$. By a comment in Section 2.2, these are

$$I_{2+3} = I_5 \qquad I_{6+7} = I_1 \qquad I_{10+11} = I_9.$$

No other reflections preserve $\underline{Hex}$ as we can see geometrically from the limited reflection symmetry of $\underline{Hex}$.

Since $H := \{T_0, T_4, T_8, I_1, I_5, I_9\}$ is a set-wise stabilizer of $\underline{Hex}$, it is a group.

From Lemma 3.1 we see that $\{T_0, T_4, T_8, I_1, I_5, I_9\}$ must also stabilize the chord collection *Hex* as a set. No other transpositions or inversions stabilize *Hex* by the argument at the outset of this proof.

(2) The only non-commutative group of order 6 is the symmetric group on three elements, denoted Sym(3), which is isomorphic to the dihedral group of order 6. The group under consideration is non-commutative, because $T_4I_1(x) = -x + 5$ while $I_1T_4(x) = -x + 3$.

□

Proposition 3.3. *The set-wise stabilizer H acts simply transitively on Hex.*

Proof. The H-orbit of $E\flat$ is all of *Hex*, as the following computation shows.



We have $|H| = 6 = |\text{orbit of } Y|$ so the Orbit-Stabilizer Theorem

$$|H| \,/\, |H_Y| = |\text{orbit of } Y|$$

implies $|H_Y| = 1$. See Proposition 2.3 2. □

Next we can investigate the subgroup of Sym(*Triads*) generated by P and L, which is called the *PL-group*.

Proposition 3.4. *The subgroup $\langle P, L\rangle$ of* Sym(*Triads*) *is dihedral of order 6.*

Proof. We first observe that P and L are involutions, that is, $P^2 = \mathrm{Id}_{Triads}$ and $L^2 = \mathrm{Id}_{Triads}$. A musical justification is the definition of "parallel" and "leading tone exchange." A mathematical justification is a direct computation of P^2 and L^2 using the formulas in (8).

Since P and L are involutions, every nontrivial element of $\langle P, L\rangle$ can be expressed as an alternating word in the letters P and L. The six functions Id_{Triads}, P, LP, PLP, $LPLP$, and $PLPLP$ are all distinct by evaluating at $E\flat$ using the following diagram from the Introduction.

(9)



From diagram (9) we also see that $(LP)^3(E\flat) = E\flat$, and for any $Y \in \{E\flat, B, G\}$, $(LP)^3(Y) = Y$. Similarly, by reading the diagram backwards (recall P and L are involutions), we see $(LP)^3(Y) = Y$ for any minor triad $Y \in \{e\flat, b, g\}$. We have similar PL-diagrams and considerations for the cycles in (4), (5), and (6), and therefore $(LP)^3 = \mathrm{Id}_{Triads}$ on the entire set *Triads* of consonant triads. Another way to see that

$(LP)^3 = \mathrm{Id}_{\mathit{Triads}}$ is to combine the observation $(LP)^3(E\flat) = E\flat$ from diagram (9) with Proposition 2.1 and the fact that *Triads* is the T/I-orbit of $E\flat$.

We next show via a word-theoretic argument that $\langle P, L\rangle$ consists only of the six functions $\mathrm{Id}_{\mathit{Triads}}$, P, LP, PLP, $LPLP$, and $PLPLP$ discussed above. From $(LP)^3 = \mathrm{Id}_{\mathit{Triads}}$, we express PL in terms of LP. Namely,

$$\begin{aligned}(LP)^3 &= \mathrm{Id}_{\mathit{Triads}}\\ (LP)^3(PL) &= (PL)\\ (LP)^2 &= PL.\end{aligned}$$

Consider any alternating word in P and L. If the right-most letter is P, then we can use $(LP)^3 = \mathrm{Id}_{\mathit{Triads}}$ to achieve an equality with one of the six functions we already have. If the right-most letter is L, then we replace each PL by $(LP)^2$ and use $L^2 = \mathrm{Id}_{\mathit{Triads}}$ if LL results on the far left. Then we have an equal function with right-most letter P, which we can then reduce to one of the six above using $(LP)^3 = \mathrm{Id}_{\mathit{Triads}}$ as we did in the first case of right-most letter P. Thus $\langle P, L\rangle = \{\mathrm{Id}_{\mathit{Triads}},\ P,\ LP,\ PLP,\ LPLP,\ PLPLP\}$.

This group is non-commutative, as $PL \neq LP$, hence it is isomorphic to Sym(3), the only non-commutative group of order 6. But Sym(3) is dihedral of order 6.

Instead of the previous paragraph, we can show $\langle P, L\rangle$ is dihedral of order 6 using a presentation. Let $t := L$ and $s := LP$, then $s^3 = e$, $t^2 = e$, and $tst = s^{-1}$. The dihedral group of order 6 is the largest group with elements s and t such that $s^3 = e$, $t^2 = e$, and $tst = s^{-1}$. But we observed from diagram (9) that $\langle P, L\rangle$ has at least six distinct elements. Hence, $\langle P, L\rangle$ is dihedral of order 6. □

Proposition 3.5. *The PL-group $\langle P, L\rangle$ acts simply transitively on Hex.*

Proof. From diagram (9) we see that $\langle P, L\rangle$ acts transitively on *Hex*. Since $\langle P, L\rangle$ and *Hex* have the same cardinality, the Orbit-Stabilizer Theorem implies that every stabilizer must be trivial. See Proposition 2.3 2. □

Lemma 3.6. *Let S be a set and suppose $G \leq \mathrm{Sym}(S)$. Suppose G acts simply transitively on an orbit $\overline{S}$, and $\overline{G}$ is the restriction of G to the orbit $\overline{S}$. Then the restriction homomorphism $G \to \overline{G}$ is an isomorphism, and $\overline{G}$ also acts simply transitively.*

Proof. Suppose $g \in G$ has restriction $\overline{g}$ with $\overline{g}\,\overline{s} = \overline{s}$ for all $\overline{s} \in \overline{S}$. Then unrestricted g has $g\overline{s} = \overline{s}$ for all $\overline{s} \in \overline{S}$, so $g = \mathrm{Id}_S$ by simple transitivity, and the kernel of the surjective homomorphism $G \to \overline{G}$ is trivial. The simple transitivity of $\overline{G}$ is fairly clear. □

Theorem 3.7 (Hexatonic Duality)**.** *The restrictions of the PL-group and $\{T_0, T_4, T_8, I_1, I_5, I_9\}$ to Hex are dual groups in* Sym(*Hex*)*, and both are dihedral of order 6.*

Proof. Let $\overline{G}$ be the restriction of the PL-group to *Hex*, and let $\overline{H}$ be the restriction of $H = \{T_0, T_4, T_8, I_1, I_5, I_9\}$ to *Hex*.

We already know that $\overline{G}$ and $\overline{H}$ are dihedral of order 6 by Propositions 3.2 and 3.4 and Lemma 3.6.

We also already know that $\overline{G}$ and $\overline{H}$ each act simply transitively on *Hex* by Propositions 3.3 and 3.4 and Lemma 3.6. We even already know that the groups $\overline{G}$ and $\overline{H}$ commute by Proposition 2.1.

We would like to conclude from the simple transitivity and commutativity of $\overline{G}$ and $\overline{H}$, using finiteness, that the centralizers in $\mathrm{Sym}(\mathit{Hex})$ are $C(\overline{G}) = \overline{H}$ and $C(\overline{H}) = \overline{G}$.

We claim that $C(\overline{G})$ acts simply transitively on *Hex*. It acts transitively, as $C(\overline{G}) \supseteq \overline{H}$ and $\overline{H}$ acts transitively. So, it suffices by Proposition 2.3 1 to prove that, for each $Y \in \mathit{Hex}$, the only element of $C(\overline{G})$ that fixes Y is the identity. Let σ be an element of $C(\overline{G})$ that fixes Y, and $\overline{g}$ any element of $\overline{G}$. Then,

$$\begin{aligned} \sigma Y &= Y \\ \overline{g}\big(\sigma Y\big) &= \overline{g}\big(Y\big) \\ \big(\overline{g}\sigma\big)Y &= \big(\overline{g}Y\big) \\ \big(\sigma\overline{g}\big)Y &= \big(\overline{g}Y\big) \\ \sigma\big(\overline{g}Y\big) &= \big(\overline{g}Y\big). \end{aligned}$$

So, not only does σ fix Y, but σ also fixes $\big(\overline{g}Y\big)$ for every $\overline{g} \in \overline{G}$. That is to say $\sigma = \mathrm{Id}_{\mathit{Hex}}$, and $C(\overline{G})$ acts simply transitively on *Hex*.

From the Orbit-Stabilizer Theorem 2.2 and the simple transitivity of the $C(\overline{G})$-action on *Hex*, we see that $C(\overline{G})$ has exactly 6 elements, see Proposition 2.3 3. But $C(\overline{G}) \supseteq \overline{H}$ and $\overline{H}$ has 6 elements, so $C(\overline{G}) = \overline{H}$.

By reversing the roles of $\overline{G}$ and $\overline{H}$ in the previous argument, we see $C(\overline{H}) = \overline{G}$. □

By the argument in the previous proof, we can also see that in the finite case, the notion of *dual groups* has a redundancy: instead of requiring the groups to centralize each other, it is sufficient to merely require that they commute.

Proposition 3.8. *Let S be a finite set. Suppose the groups $G, H \leq \mathrm{Sym}(S)$ act simply transitively on S, and that G and H commute in the sense that $gh = hg$ for all $g \in G$ and $h \in H$. Then $C(G) = H$ and $C(H) = G$.*

Proof. This follows from the proof of Theorem 3.7, when we replace $\overline{G}$ and $\overline{H}$ by G and H, and ignore the specifics of *Hex* and the group structures. □

Remark 3.9 (Comparison with the Proof Strategy of Crans–Fiore–Satyendra in [4])**.** There are several differences between the proof strategy of Hexatonic Duality in present Theorem 3.7 and the proof strategy of T/I-PLR duality in Theorem 6.1 of [4]. In the present paper, we first proved that the concerned groups act simply transitively, and determined their structure, and only then showed that the groups exactly centralize each other. In [4], on the other hand, the determination of the size of the PLR-group was postponed until after the centralizer $C(T/I)$ was seen to act simply, i.e. that each stabilizer $C(T/I)_Y$ is trivial. Then, from these trivial stabilizers, the Orbit-Stabilizer Theorem, the earlier observation that $24 \leq |PLR\text{-group}|$, and the consequence

$$24 \leq |PLR\text{-group}| \leq |C(T/I)| \leq |\text{orbit of } Y| \leq 24$$

on page 492, the authors of [4] simultaneously conclude that the PLR-group has 24 elements and is the centralizer of T/I.

A slight simplification of the aforementioned inequality would be an argument like the one in the present paper: observe that the PLR-group acts transitively on the

24 consonant triads because of the Cohn LR-sequence (recalled on page 487 of [4]), then $C(T/I)$ must act transitively as it contains the PLR-group, and then the Orbit-Stabilizer Theorem and the trivial stabilizers imply that $|C(T/I)|$ must be 24, so the PLR-group also has 24 elements. Also, instead of postponing the proof that the PLR-group has exactly 24 elements from Theorem 5.1 of [4] until the aforementioned inequality in Theorem 6.1, one could do a word-theoretic argument in Theorem 5.1 to see that the PLR-group has exactly 24 elements, similar to the present argument in Proposition 3.4.

Remark 3.10. For an explicit computation of the four hexatonic cycles as orbits of the PL-group, see Oshita [24], which was also an undergraduate research project with the second author of the present article. The preprint [24] includes a sketch that $\langle P, L\rangle \cong \mathrm{Sym}(3)$.

Remark 3.11 (Alternative Derivation using the Sub Dual Group Theorem)**.** Hexatonic Duality Theorem 3.7 can also be proved using the Sub Dual Group Theorem 3.1 of Fiore–Noll in [6], *if one assumes already* the duality of the T/I-group and PLR-group (maximal smoothness is not discussed in [6]). Fiore–Noll apply the Sub Dual Group Theorem to the construction of dual groups on the hexatonic cycles in [6, Section 3.1]. The method is to select G_0 to be the PL-group, select $s_0 = E\flat$, and compute $S_0 := G_0 s_0 = \mathit{Hex}$, and then the dual group will consist of the restriction of those elements of the T/I-group that map $E\flat$ into S_0.

Notice that in the present paper, on the other hand, we *first* determined which transpositions and inversions preserve *Hex* in Proposition 3.2, and then proved duality, whereas the application of the Sub Dual Group Theorem of Fiore–Noll starts with the PL-group and determines from it the dual group as (the restrictions of) those elements of the T/I-group that map $E\flat$ into S_0. Notice also, in the present paper we determined that the PL-group and its dual H are dihedral of order 6, but that the Sub Dual Group Theorem of Fiore–Noll does not specify which group structure is present. In any case, Clampitt explicitly wrote down all 6 elements of each group in permutation cycle notation in [2].

The present paper is complementary to the work [6] of Fiore–Noll in that we work very closely with the specific details of the groups and sets involved to determine one pair of dual groups in an illustrative way, rather than appealing to a computationally and conceptually convenient theorem. Fiore–Noll however also use their Corollary 3.3 to compute the other hexatonic duals via conjugation, as summarized in Table 2.

The application of the Sub Dual Group Theorem to construct dual groups on octatonic systems is also treated in [6], and utilized in [8].

Remark 3.12 (Other Sources on Group Actions)**.** Music theoretical group actions on chords have been considered by many, many authors over the past century. In addition to the selected references of Babitt, Forte, and Morris above, we also mention the expansive and influential work of Mazzola and numerous collaborators [15, 16, 17].

Remark 3.13 (Discussion of Local Diatonic Containment of Hexatonic Cycles)**.** No hexatonic cycle is contained entirely in a single diatonic set, as one can see from any of the cycles (3)–(6). However, one can consider a sequence of diatonic sets that changes

k	$kHex$	kHk^{-1} = dual group to PL-group on $kHex$
Id_{Triads}	$\{E\flat, e\flat, B, b, G, g\}$	$H = \{T_0, T_4, T_8, I_1, I_5, I_9\}$
T_1	$\{E, e, C, c, A\flat, a\flat\}$	$\{T_0, T_4, T_8, I_3, I_7, I_{11}\}$
T_2	$\{F, f, C\sharp, c\sharp, A, a\}$	$\{T_0, T_4, T_8, I_5, I_9, I_1\}$
T_3	$\{F\sharp, f\sharp, D, d, B\flat, b\flat\}$	$\{T_0, T_4, T_8, I_7, I_{11}, I_3\}$

TABLE 2. The 4 hexatonic cycles as PL-orbits and the respective dual groups determined as conjugations of H via the Sub Dual Group Theorem of Fiore–Noll.

along with the hexatonic cycle and contains each respective triad, as Douthett does in [5, Table 4.7]. After transposing and reversing Douthett's table, we see a sequence of diatonic sets such that each diatonic set contains the respective triad of (3).

Triad	$E\flat$	$e\flat$	B	b	G	g
In Scale	$E\flat$-major	$D\flat$-major	B-major	A-major	G-major	F-major

This sequence of diatonic sets (indicated via major scales) descends by a whole step each time, so is as evenly distributed as possible.

Other diatonic set sequences also contain the hexatonic cycle, though unfortunately there is no maximally smooth cycle of diatonic sets that does the job (recall that the diatonic sets can only form a cycle of length 12). But it is possible to have a maximally smooth sequence of diatonic sets that covers four hexatonic triads. We list all possible diatonic sets containing the respective hexatonic chords.[1]

Triad	$E\flat$	$e\flat$	B	b	G	g
In Major Scales	$E\flat$	$G\flat$	B	D	G	$A\flat$
	$B\flat$	$\mathbf{D}\flat$	$F\sharp$	$\mathbf{A}$	D	$\mathbf{F}$
	$\mathbf{A}\flat$	B	$\mathbf{E}$	G	$\mathbf{C}$	$E\flat$

Double vertical dividing lines indicate maximally smooth transitions between consecutive diatonic sets. As indicated by double vertical dividing lines, the transition from a minor triad to its consecutive major in a hexatonic cycle via L is contained in three maximally smooth transitions of diatonic sets. On the other hand, the transition from a major triad to its consecutive minor in a hexatonic cycle via P is contained in only one maximally smooth transition of diatonic sets, as indicated by the bold letters. Altogether, we can trace three maximally smooth chains of four major scales that contain part of the hexatonic cycle (3).

$$B - E - A - D$$

[1]Recall that major chords only occur with roots on major scale degrees 1, 4, and 5, so we determine in the table the scales containing a given major triad by considering the root, a perfect fourth below the root, and a perfect fifth below the root. Minor scales can only occur with roots on major scale degrees 2, 3, and 6, so we determine in the table the scales containing a given minor triad by considering a major sixth below the root, a whole step below the root, and a major third below the root. This nonconsistent major/minor ordering allows us to see (at double vertical dividing lines) all three maximally smooth transitions from diatonic sets containing a given a minor triad to a diatonic set containing its consecutive major in a hexatonic cycle.

$$G - C - F - B\flat$$

$$E\flat - A\flat - D\flat - F\sharp$$

Local containment of hexatonic cycles in diatonic chains has ramifications for music analysis. Jason Yust proposes in [26] and [27] to include diatonic contexts into analyses involving PL-cycles or PR-cycles, and he provides analytical tools to do so. We thank Thomas Noll for proposing and discussing this extended Remark 3.13 with us.

4. Conclusion

We began this article with Cohn's proposal that the maximal smoothness of consonant triads is a key factor for their privileged status in late-nineteenth century music. Indeed, consonant triads and their complements are the only tone collections that accommodate short maximally smooth cycles. The four maximally smooth cycles of consonant triads, the so-called hexatonic cycles of Cohn, can be described transformationally as alternating applications of the neo-Riemannian "parallel" and "leading tone exchange" transformations. Cohn interpreted Wagner's Grail motive in terms of a cyclic group action on the hexatonic cycle *Hex*, whereas Clampitt used the PL-group and the transposition-inversion subgroup we called H in Proposition 3.2. In the present article, we proved the Lewinian duality between these latter two groups, which was discussed by Clampitt in [2].

For perspective, we mention that simply transitive group actions correspond to the *generalized interval systems* of Lewin, see the very influential original source [14], or see [8, Section 2] for an explanation of some aspects. Dual groups correspond to dual generalized interval systems: the transpositions of one system are the interval preserving bijections of the other. Clampitt [2] explained the coherent perceptual basis of the three generalized interval systems associated to the three group actions on *Hex* by Cohn's cyclic group, the PL-group, and the H group. He employed the coherence of generalized interval systems to incorporate the final $D\flat$ of the Grail motive into his interpretation via a conjugation-modulation of a subsystem.

Acknowledgements

This paper is the extension of an undergraduate research project of student Cameron Berry with Professor Thomas Fiore at the University of Michigan-Dearborn. Cameron Berry thanks Professor Thomas Fiore for all of the time he spent assisting and encouraging him during Winter semester 2014. We both thank Mahesh Agarwal for a suggestion to use a word-theoretic argument in Theorem 5.1 of [4], which we implemented in the present Proposition 3.4. We thank Thomas Noll for proposing and discussing Remark 3.13. Thomas Fiore thanks Ramon Satyendra for introducing him to hexatonic cycles and generalized interval systems back in 2004.

Thomas Fiore was supported by a Rackham Faculty Research Grant of the University of Michigan. He also thanks the Humboldt Foundation for support during his 2015-2016 sabbatical at the Universität Regensburg, which was sponsored by a Humboldt Research Fellowship for Experienced Researchers. Significant progress on this project was completed during the fellowship.

References

[1] Milton Babbitt. Some aspects of twelve-tone composition. *The Score and IMA Magazine*, 12:53–61, 1955.

[2] David Clampitt. Alternative interpretations of some measures from Parsifal. *Journal of Music Theory*, 42(2):321–334, 1998.

[3] Richard Cohn. Maximally smooth cycles, hexatonic systems, and the analysis of late-romantic triadic progressions. *Music Analysis*, 15(1):9–40, 1996.

[4] Alissa S. Crans, Thomas M. Fiore, and Ramon Satyendra. Musical actions of dihedral groups. *American Mathematical Monthly*, 116(6):479–495, 2009.

[5] Jack Douthett. Filtered point-symmetry and dynamical voice leading. In Jack Douthett, Martha M. Hyde, and Charles J. Smith, editors, *Music Theory and Mathematics: Chords, Collections, and Transformations*, volume 50 of *Eastman Studies in Music*. University of Rochester Press, 2008.

[6] Thomas M. Fiore and Thomas Noll. Commuting groups and the topos of triads. In Carlos Agon, Emmanuel Amiot, Moreno Andreatta, Gerard Assayag, Jean Bresson, and John Mandereau, editors, *Proceedings of the 3rd International Conference Mathematics and Computation in Music - MCM 2011*, Lecture Notes in Computer Science. Springer, 2011.

[7] Thomas M. Fiore, Thomas Noll, and Ramon Satyendra. Incorporating voice permutations into the theory of neo-Riemannian groups and Lewinian duality. In *Mathematics and Computation in Music*, volume 7937 of *Lecture Notes in Compuer Science*, pages 100–114. Springer, Heidelberg, 2013.

[8] Thomas M. Fiore, Thomas Noll, and Ramon Satyendra. Morphisms of generalized interval systems and PR-groups. *Journal of Mathematics & Music*, 7(1):3–27, 2013.

[9] Thomas M. Fiore and Ramon Satyendra. Generalized contextual groups. *Music Theory Online*, 11(3), 2005, `http://mto.societymusictheory.org`.

[10] Allen Forte. *The Structure of Atonal Music*. Yale University Press, New Haven, CT, 1973.

[11] Harald Fripertinger and Peter Lackner. Tone rows and tropes. *Journal of Mathematics and Music*, 9(2):111–172, 2015.

[12] Julian Hook. Why are there twenty-nine tetrachords? A tutorial on combinatorics and enumeration in music theory. *Music Theory Online*, 13(4), 2007.

[13] Julian Hook and Robert Peck. Introduction to the special issue on tone rows and tropes. *Journal of Mathematics and Music*, 9(2):109–110, 2015.

[14] David Lewin. *Generalized Musical Intervals and Transformations*. Yale University Press, New Haven, 1987.

[15] G. Mazzola. *Gruppen und Kategorien in der Musik*, volume 10 of *Research and Exposition in Mathematics*.

[16] Guerino Mazzola. *Geometrie der Töne*. Birkhäuser Verlag, Basel, 1990.

[17] Guerino Mazzola. *The topos of music*. Birkhäuser Verlag, Basel, 2002. Geometric logic of concepts, theory, and performance, in collaboration with Stefan Göller and Stefan Müller.

[18] Brian J. McCartin. Prelude to musical geometry. *The College Mathematics Journal*, 29:354–370, 1998.

[19] Andrew Mead. Remarks on "Tone Rows and Tropes" by Harald Fripertinger and Peter Lackner. *Journal of Mathematics and Music*, 9(2):173–178, 2015.

[20] Robert Morris. *Composition with Pitch Classes: A Theory of Compositional Design*. Yale University Press, New Haven and London, 1987.

[21] Robert Morris. *Class Notes for Atonal Music Theory*. Frog Peak Music, 1991.

[22] Robert Morris. *Class Notes for Advanced Atonal Music Theory*. Frog Peak Music, 2001.

[23] Robert Morris. Review of "Tone Rows and Tropes" by Harald Fripertinger and Peter Lackner. *Journal of Mathematics and Music*, 9(2):179–195, 2015.

[24] Kenneth Oshita. The hexatonic systems under neo-Riemannian theory: an exploration of the mathematical analysis of music. *REU Project, University of Chicago*, 2009, http://www.math.uchicago.edu/~may/VIGRE/VIGRE2009/REUPapers/Oshita.pdf.
[25] John Rahn. *Basic Atonal Theory*. Schirmer Books, 1987.
[26] Jason Yust. Tonal prisms: Iterated quantization in chromatic tonality and ravel's ondine. *Journal of Mathematics and Music*, 7(2):145–165, 2013.
[27] Jason Yust. Distorted continuity: Chromatic harmony, uniform sequences, and quantized voice leadings. *Music Theory Spectrum*, 37(1):120–143, 2015.

Department of Mathematics, Michigan State University, 619 Red Cedar Road, East Lansing, MI 48824
E-mail address: berrycam@msu.edu

Thomas M. Fiore, Department of Mathematics and Statistics, University of Michigan-Dearborn, 4901 Evergreen Road, Dearborn, MI 48128
and
NWF I - Mathematik, Universität Regensburg, Universitätsstrasse 31, 93040 Regensburg, Germany
E-mail address: tmfiore@umich.edu
URL: http://www-personal.umd.umich.edu/~tmfiore/